

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search PubMed

for

Go

Clear

☒ Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Pigment Cell Res. 1993 Dec;6(6):394-9.

Related Articles, L

Improved tyrosinase activity stains in polyacrylamide electrophoresis gels.

Jimenez-Cervantes C, Valverde P, Garcia-Borron JC, Solano F, Lozano JA.

Department of Biochemistry, Molecular Biology and Immunology, School of Medicine, University of Murcia, Spain.

Mammalian tyrosinase exists in a variety of subcellular locations and maturation states that result from a complex post-translational processing with possible regulatory implications. So far, SDS-PAGE has proven to be the method of choice for the resolution of tyrosinase isoforms. However, the relatively poor sensitivity of the currently available specific activity stain based on incubation of the gels with L-dopa until the formation of melanin has severely limited the use of electrophoresis in regulation studies. Two alternative staining procedures are presented and discussed. The first one involves the fluorographic detection of radioactive melanin after incubation of the gels in the presence of L-[3-¹⁴C]-dopa. A similar method has already been used by others (Tsukamoto et al., 1992, Pigment Cell Res. [Suppl.] 2:84-89), but its performance has not yet been compared to the one of the dopa procedure. The sensitivity of this method can be varied by adjusting the isotopic dilution of the tracer and/or the time of exposure of the gel, but it is at least ten times higher than the one of the colorimetric stain. Moreover, the intensity of the bands is proportional to the initial tyrosinase activity over a wide range. Using this procedure, the activity present in the different subcellular fractions of melanocytes in culture can be easily detected. The second procedure involves the formation of a colored adduct between dopaquinone and MBTH. (ABSTRACT TRUNCATED AT 250 WORDS)

PMID: 7511805 [PubMed - indexed for MEDLINE]

Display

Abstract


Show:

20

Sort

Send to

Text



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Bc

Search PubMed for

About Entrez

Limits Preview/Index History Clipboard Details

Abstract Text

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

☐ 1: Int J Hyperthermia. 2002 Nov-Dec;18(6):563-75.

Related Articles, L

MetaPress

Heat shock protein 70: role in antigen presentation and immune stimulation.

Milani V, Noessner E, Ghose S, Kuppner M, Ahrens B, Scharner A, Gastpar R, Issels RD.

KKG Hyperthermie, GSF-National Research Center for Environment and Health, 81377 Munich, Germany.

Heat shock proteins (HSP) when released into the extracellular milieu can act simultaneously as a source of antigen due to their ability to chaperone peptid and as a maturation signal for dendritic cells, thereby inducing DCs to cross-present antigens to CD8+ T-cells. HSP can also act independently from associated peptides, stimulating the innate immune system. Previous results regarding the activation of NK cells by HSP70 cell surface expression on tumour cells and soluble HSP70 will be further covered elsewhere within this issue. For cross-presentation, HSP70-peptide complexes (HSP70-PC) were used from two human melanoma cell lines that differ in the expression of the tumour-associated antigen tyrosinase. Purified HSP70-PC consists of both the constitutively expressed HSC70 and the inducible HSP70. HSP70-peptide complexes purified from tyrosinase positive (HSP70-PC/tyr+) human melanoma cells, incubated with immature DCs, results in the activation of HLA-A*0201-restricted tyrosinase peptide-specific T-cells. Receptor-mediated uptake of HSP70-PC by DCs and intracellular transport are required for efficient MHC class I restricted cross-presentation of chaperoned peptides. Demonstration of HSP70-PC mediated cross-presentation of such non-mutated naturally expressed tumour antigens is of special clinical interest with regard hyperthermia. Tumour regression and improved local control have been shown within clinical phase II/III trials integrating regional hyperthermia combined with radiation and/or chemotherapy in multimodal treatment strategies. According to the proposed concept, local necrosis induced by hyperthermic treatment induces the release of HSPs, followed by uptake, processing and presentation of associated peptides by DCs. By acting as chaperone and a signal for DC maturation, HSP70-PC might efficiently prime circulating T-cells. Therefore, upregulating HSP70 and causing local necrosis in tumour tissue by hyperthermia offers great potential as a new approach to directly activate the immune system.